

Editorial

Lung-to-head ratio: a need to unify the technique

J. C. JANI*†, C. F. A. PERALTA‡ and
K. H. NICOLAIDES§¶

†Fetal Medicine Unit of the University Hospital Brugmann, Brussels, Belgium; ‡Department of Obstetrics and Gynecology, Center for Integral Assistance to Women's Health, State University of Campinas Medical School, Campinas, Brazil; §Harris Birthright Research Centre of Fetal Medicine, King's College Hospital, London, UK; ¶Department of Fetal Medicine, University College London NHS Foundation Trust, London, UK

*Correspondence: Dr J. Jani, CHU Brugmann, Department of Obstetrics and Gynecology, Place A. Van Gehuchten n°4, 1020 Bruxelles, Belgium (e-mail: jackjani@hotmail.com)

Congenital diaphragmatic hernia (CDH) is associated with a high postnatal mortality rate due to pulmonary hypoplasia and/or hypertension. Antenatal prediction of postnatal outcome is based essentially on the assessment of the degree of intrathoracic compression of the lungs, and the most widely used method is ultrasonographic measurement of the fetal lung area-to-head circumference ratio (LHR).

In the assessment of fetuses with CDH and the counseling of parents as to the available options (including endoscopic occlusion of the fetal trachea^{1–3}, expectant management and pregnancy termination) for subsequent management of the pregnancy, it is imperative that prediction of outcome is as accurate as possible. Consequently, operators undertaking measurement of LHR should be trained appropriately⁴ and correct algorithms used for interpretation of the results.

Evolution in the application of LHR

In the first description of this technique, the area of the right lung in fetuses with left-sided CDH and intrathoracic herniation of the liver was measured at ≤ 25 weeks' gestation⁵. The right lung area was estimated, in the cross-sectional plane of the thorax used for examination of the four-chamber view of the heart, by multiplication of the longest diameter of the lung by its longest perpendicular diameter (Figure 1a)⁵. However, in the figure provided in that article to explain the technique, estimation of the lung area was obtained by multiplication of the anteroposterior diameter of the lung at the mid-clavicular line by the perpendicular diameter at the midpoint of the anteroposterior diameter (Figure 1b). The authors of that study recognized that the lung area increases

with gestational age and attempted to create a gestation-independent measurement by dividing the lung area by the head circumference.

Subsequent studies utilizing LHR in the prediction of postnatal outcome in fetuses with CDH have generally confirmed that survival improves with increasing LHR^{5–18}. However, there are large differences in reported results, possibly because of the wide range in gestational age, from 16–39 weeks, at which the LHR was measured, the small number of patients examined in most studies and the extension of the use of LHR from cases with left-sided CDH with intrathoracic liver to those with right-sided defects and those without liver herniation. Additionally, although all studies used the same reference to explain the method of measuring lung area, i.e. multiplication of the longest diameter of the lung by its longest perpendicular diameter⁵, in reality there were inconsistencies in the figures they provided to explain their technique.

A cross-sectional study of 650 normal singleton pregnancies at 12–32 weeks of gestation established reference intervals for the right and left lung areas and LHRs¹⁹. The lung areas were measured, on the cross-sectional plane of the thorax used for examination of the four-chamber view of the heart, by three different techniques (Figures 1a–c): firstly, multiplication of the longest diameter of the lung by its longest perpendicular diameter; secondly, multiplication of the anteroposterior diameter of the lung at the mid-clavicular line by the perpendicular diameter at the midpoint of the anteroposterior diameter; and thirdly, manual tracing of the limits of the lungs. The data of this study demonstrated that between 12 and 32 weeks of gestation there is an 18-fold increase in lung area and a four-fold increase in head circumference. Consequently, the left and right LHR increase exponentially with gestation (Figure 2). The most reproducible way of measuring the lung area was by manual tracing of the limits of the lungs and the least reproducible was by multiplying the longest diameter of the lungs by their longest perpendicular diameter. Furthermore, the method employing the longest diameter, compared with the tracing method, overestimated both left and right lung areas by about 45% and the method employing the anteroposterior diameter overestimated the area of the right lung by about 35%, but not that of the left lung.

A subsequent study of 354 fetuses with isolated CDH, reported that the LHR at 18–38 weeks' gestation increased with gestational age, having estimated the lung area by multiplication of the longest diameter of the lung by its longest perpendicular diameter²⁰. The authors proposed the introduction of a new measurement, the observed to expected (o/e) LHR, which was independent of gestational age. This method is applicable to both fetuses with left-sided CDH, with or without intrathoracic




Figure 1 Ultrasound images in fetuses with normal lungs (a–c) and left-sided congenital diaphragmatic hernia (d–f) at the level of the four-chamber view of the heart. Three methods of measuring the lung area are illustrated: (a,d) multiplication of the longest diameter of the lung by its longest perpendicular diameter (longest diameter method); (b,e) multiplication of the anteroposterior diameter of the lung at the mid-clavicular line by the perpendicular diameter at the midpoint of the anteroposterior diameter (anteroposterior method); and (c,f) tracing of the limits of the lungs (tracing method).




Figure 2 Right (a) and left (b) lung areas, by the longest diameter method, to head circumference ratio (LHR) according to gestational age in normal fetuses.




Figure 3 Observed to expected right lung area-to-head circumference ratio (LHR) according to gestational age in normal fetuses (shaded area represents 95% reference range) and in individual fetuses with left-sided congenital diaphragmatic hernia (circles with regression lines), showing fetuses that died (—●—) and survivors (---○---).

herniation of the liver, and fetuses with right-sided CDH. Furthermore, in both left- and right-sided CDH, measurement of the o/e LHR was useful in the prediction of subsequent survival (Figure 3)²⁰. In fetuses with left-sided CDH the survival rate was substantially lower in those with intrathoracic herniation of the liver than in those with the liver confined to the abdomen (53% vs 75%). However, logistic regression analysis demonstrated that the position of the liver did not provide significant contribution independent of the o/e LHR in the prediction of survival²⁰.

Protocol for measurement of LHR

The following criteria should be applied in the measurement of LHR.

1. An axial view at the level of the four-chamber view of the heart should be obtained.

2. The view should be such that the lung contralateral to the CDH is closer to the probe than the herniated structures.
3. Care should be taken to avoid shadows produced by the ribs on the lung to be measured, by placing the transducer so that the ultrasound beam is parallel to the ribs and crosses the intercostal space.
4. The image should be frozen prior to final magnification to ensure that all landmarks are clearly visible. The image should then be magnified so that the axial view of the fetal thorax occupies the whole screen.
5. The calipers should be placed according to the method being used to measure the lung area: longest diameter, anteroposterior diameter or tracing method (Figures 1a–c). In all methods, the area of the lung (in square millimeters) is divided by the head circumference (in millimeters) to obtain the LHR. For all methods, care should be taken not to include the myocardium in the measurement of the lung area. The preferred method is the tracing one since it is the most reproducible¹⁹, especially when the lung contralateral to the CDH has an unusual and irregular shape (Figure 4).
6. When the measurement of LHR is completed, it should be divided by the expected mean LHR for gestational age, so that the o/e LHR is obtained. Table 1 provides the formulae for the expected mean LHR for right and left lungs using the three different methods of measuring the lung area.
7. The minimum number of LHR measurements required to achieve competence in the technique is 70⁴. Training should be done under the supervision of a sonographer with sufficient expertise in measuring LHR. An alternative is to send the images to a center with experts in LHR measurement for evaluation. It is acceptable that part of the training is performed on fetuses with normal lungs.

Limitations of LHR in the prediction of outcome

The measurement of fetal LHR has been used for the prediction of pulmonary hypoplasia and/or hypertension in the neonatal period. Studies have reported a high association between o/e LHR and such outcome both in fetuses managed expectantly and in those treated by fetoscopic occlusion of the trachea^{2,20}. However, postnatal survival is related not only to the degree of fetal lung development but also to the expertise of the neonatal center caring for the affected infant. Consequently, it is necessary to adjust the o/e LHR according to the level of neonatal care.

Table 1 Formulae for expected fetal lung area-to-head circumference ratio (LHR) in left- and right-sided congenital diaphragmatic hernia (CDH), using different methods for measurement of the lung area, based on gestational age (GA) in weeks with decimals

Lung area measurement	Right LHR in left CDH	Left LHR in right CDH
Longest diameter method	$-3.4802 + (0.3995 \times GA) - (0.0048 \times GA^2)$	$-2.5957 + (0.3043 \times GA) - (0.0042 \times GA^2)$
Anteroposterior diameter method	$-3.1597 + (0.3615 \times GA) - (0.0041 \times GA^2)$	$-1.0224 + (0.1314 \times GA) - (0.0011 \times GA^2)$
Tracing method	$-2.3271 + (0.27 \times GA) - (0.0032 \times GA^2)$	$-1.4994 + (0.1778 \times GA) - (0.0021 \times GA^2)$




Figure 4 Ultrasound images at the level of the four-chamber view of the heart in a fetus with left-sided congenital diaphragmatic hernia, demonstrating some of the difficulties in estimation of the lung area. In (a) there is extension of the contralateral lung around the heart (arrow), making it difficult to decide where to place the calipers for measurement of the longest diameter; in (b) and (c) the stomach and spleen, respectively, distort the shape of the lung, making it difficult to decide where to place the calipers for measurement of the transverse diameter.

Magnetic resonance imaging (MRI) provides more accurate measurement of lung volumes than does sonographic assessment by LHR²¹. Additionally, measurements obtained by MRI are easier to standardize and allow assessment of both lungs. However, MRI is confined to specialist centers and therefore measurement of LHR will, in the foreseeable future, remain the first-line method of assessment of the severity of CDH. Ultimately, if ongoing trials demonstrate that fetoscopic occlusion of the fetal trachea is beneficial, an increasing number of affected fetuses could be managed in specialist centers where the main method of assessment of the severity of CDH would be MRI.

REFERENCES

- Deprest J, Gratacos E, Nicolaides KH; FETO Task Group. Fetoscopic tracheal occlusion (FETO) for severe congenital diaphragmatic hernia: evolution of a technique and preliminary results. *Ultrasound Obstet Gynecol* 2004; **24**: 121–126.
- Jani JC, Nicolaides KH, Gratacos E, Valencia CM, Doné E, Martinez JM, Gucciardo L, Cruz R, Deprest JA. Severe diaphragmatic hernia treated by fetal endoscopic tracheal occlusion. *Ultrasound Obstet Gynecol* 2009; **34**: 304–310.
- Ruano R, Yoshisaki CT, da Silva MM, Ceccon ME, Grasi MS, Tannuri U, Zugaib M. A randomized controlled trial of fetal endoscopic tracheal occlusion versus postnatal management of severe isolated congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2012; **39**: 20–27.
- Cruz-Martinez R, Figueras F, Moreno-Alvarez O, Gomez O, Martinez JM, Hernandez-Andrade E, Gratacos E. Learning curve for the lung area to head circumference ratio measurement in fetuses with congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2010; **36**: 32–36.
- Metkus AP, Filly RA, Stringer MD, Harrison MR, Adzick NS. Sonographic predictor of survival in fetal diaphragmatic hernia. *J Pediatr Surg* 1996; **31**: 148–151.
- Lipshutz GS, Albanese CT, Feldstein VA, Jennings RW, Housley HT, Beech R, Farrell JA, Harrison MR. Prospective analysis of lung-to-head ratio predicts survival for patients with prenatally diagnosed congenital diaphragmatic hernia. *J Pediatric Surg* 1997; **32**: 1634–1636.
- Harrison MR, Mychaliska GB, Albanese CT, Jennings RW, Farrell JA, Hawgood S, Sandberg P, Levine AH, Lobo E, Filly RA. Correction of congenital diaphragmatic hernia in utero IX: Fetuses with poor prognosis (liver herniation and low lung-to-head ratio) can be saved by fetoscopic temporary tracheal occlusion. *J Pediatr Surg* 1998; **33**: 1017–1023.
- Flake AW, Crombleholme TM, Johnson MP, Howell LJ, Adzick NS. Treatment of severe congenital diaphragmatic hernia by fetal tracheal occlusion: Clinical experience with fifteen cases. *Am J Obstet Gynecol* 2000; **183**: 1059–1066.
- Sbragia L, Paek B, Filly RA, Harrison MR, Farrell J, Farmer D, Albanese CT. Congenital diaphragmatic hernia without herniation of the liver: Does the lung-to-head ratio predict survival? *J Ultrasound Med* 2000; **19**: 845–848.
- Laudy JAM, Van Gucht M, Van Dooren MF, Wladimiroff JW, Tibboel D. Congenital diaphragmatic hernia: an evaluation of the prognostic value of the lung-to-head ratio and other prenatal parameters. *Prenat Diagn* 2003; **23**: 634–639.
- Heling KS, Wauer RR, Hammer H, Bollmann R, Chaoui R. Reliability of the lung-to-head ratio in predicting outcome and neonatal ventilation parameters in fetus with congenital diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2005; **25**: 112–118.
- Jani J, Keller RL, Benachi A, Nicolaides KH, Favre R, Gratacos E, Laudy J, Eisenberg V, Eggink A, Vaast P, Deprest J.

- Prenatal prediction of survival in isolated left-sided diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2006; **27**: 18–22.
13. Arkovitz MS, Russo M, Devine P, Budhorick N, Stolar CJ. Fetal lung-head ratio is not related to outcome for antenatal diagnosed congenital diaphragmatic hernia. *J Pediatr Surg* 2007; **42**: 107–110.
 14. Hedrick HL, Danzer E, Merchant A, Bebbington MW, Zhao H, Flake AW, Johnson MP, Liechty KW, Howell LJ, Wilson RD, Adzick NS. Liver position and lung-to-head ratio for prediction of extracorporeal membrane oxygenation and survival in isolated left congenital diaphragmatic hernia. *Am J Obstet Gynecol* 2007; **197**: 422.e1–4.
 15. Yang SH, Nobuhara KK, Keller RL, Ball RH, Goldstein RB, Feldstein VA, Callen PW, Filly RA, Farmer DL, Harrison MR, Lee H. Reliability of the lung-to-head ratio as a predictor of outcome in fetuses with isolated left congenital diaphragmatic hernia at gestation outside 24–26 weeks. *Am J Obstet Gynecol* 2007; **197**: 30.e1–7.
 16. Datin-Dorriere V, Rouzies S, Taupin P, Walter-Nicolet E, Benachi A, Sonigo P, Mitanchez D. Prenatal prognosis in isolated congenital diaphragmatic hernia. *Am J Obstet Gynecol* 2008; **198**: 80.e1–5.
 17. Alfaraj MA, Shah PS, Bohn D, Pantazi S, O'Brien K, Chiu PP, Gaiteiro R, Ryan G. Congenital diaphragmatic hernia: lung-to-head ratio and lung volume for prediction of outcome. *Am J Obstet Gynecol* 2011; **205**: 43.e1–8.
 18. Aspelund G, Fisher JC, Simpson LL, Stolar CJ. Prenatal lung-head ratio: threshold to predict outcome for congenital diaphragmatic hernia. *J Matern Fetal Neonatal Med* 2011; Oct 4. [Epub ahead of print].
 19. Peralta CFA, Cavoretto P, Csapo B, Vandecruys H, Nicolaides KH. Assessment of lung area in normal fetuses at 12–32 weeks. *Ultrasound Obstet Gynecol* 2005; **26**: 718–724.
 20. Jani J, Nicolaides KH, Keller RL, Benachi A, Peralta CFA, Favre R, Moreno O, Tibboel D, Lipitz S, Eggink A, Vaast P, Allegaert K, Harrison M, Deprest J, on behalf of the Antenatal-CDH-Registry Group. Observed to expected lung area to head circumference ratio in the prediction of survival in fetuses with isolated diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2007; **30**: 67–71.
 21. Jani J, Cannie M, Sonigo P, Robert Y, Moreno O, Benachi A, Vaast P, Gratacos E, Nicolaides KH, Deprest J. Value of prenatal magnetic resonance imaging in the prediction of postnatal outcome in fetuses with diaphragmatic hernia. *Ultrasound Obstet Gynecol* 2008; **32**: 793–799.